




The Sarawak Museum Journal
Vol. LXXIX No. 100
December 2018



ISSN: 0375-3050

Citation: *Sarawak Museum Journal*, LXXIX (100) (2018): 287-294

CAMERA TRAPPING: A TOOL TO STUDY HORNBILLS?

Thaqifah S. Jailan, Melynda Cheok Ka Yi, Sally S. Kaicheen, Hidayah
Nurul-Asna, Philoveny Pengiran and Jayasilan Mohd-Azlan*

Animal Resource Science and Management Programme, Faculty of Resource Science and
Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak

*Corresponding author. Email: azlan@unimas.my

ABSTRACT

Of the 57 species of hornbills in the world, 25 species are recorded in Africa, 32 species are recorded in Asia but only eight occur in Borneo. However, due to logging, hunting and forest fragmentation, the Bornean lowland forest often does not host all the species at once. The study of using remotely triggered infrared cameras in avifauna research is commonly used for medium to large terrestrial mammals, and birds, such as pheasants. Documentation of rare and new species presence records, behavioural studies, and activity patterns could also be included in the application of camera trapping in the study of birds. From the 304 camera trapping sites distributed throughout Sarawak, two species of hornbills were photographed during the study period: the White-Crowned Hornbill *Berenicornis comatus* and Asian Black Hornbill *Anthracoceros malayanus*. These images represent approximately 0.01% (34 out of 29,618 images) of the overall independent photos from a total effort of 29,470 camera trapping nights. Although data are insufficient to support the activity overlap between these two species, some baseline information on the behaviour of hornbills has been observed.

Keywords: activity overlap, Borneo, camera trapping, hornbills

CAMERA TRAPPING: A TOOL TO STUDY HORNBILLS?

Thaqifah S. Jailan, Melynda Cheok Ka Yi, Sally S. Kaicheen, Hidayah
Nurul-Asna, Philoveny Pengiran and Jayasilan Mohd-Azlan*

*Animal Resource Science and Management Programme, Faculty of Resource Science and
Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak*

**Corresponding author. Email: azlan@unimas.my*

Abstract

Of the 57 species of hornbills in the world, 25 species are recorded in Africa, 32 species are recorded in Asia but only eight occur in Borneo. However, due to logging, hunting and forest fragmentation, the Bornean lowland forest often does not host all the species at once. The study of using remotely triggered infrared cameras in avifauna research is commonly used for medium to large terrestrial mammals, and birds, such as pheasants. Documentation of rare and new species presence records, behavioural studies, and activity patterns could also be included in the application of camera trapping in the study of birds. From the 304 camera trapping sites distributed throughout Sarawak, two species of hornbills were photographed during the study period: the White-Crowned Hornbill *Berenicornis comatus* and Asian Black Hornbill *Anthracoceros malayanus*. These images represent approximately 0.01% (34 out of 29,618 images) of the overall independent photos from a total effort of 29,470 camera trapping nights. Although data are insufficient to support the activity overlap between these two species, some baseline information on the behaviour of hornbills has been observed.

Keywords: activity overlap, Borneo, camera trapping, hornbills

INTRODUCTION

Hornbills in Borneo are canopy-dwelling, monogamous pairs and hole-nesters (Kinnaird & O'Brien 2007; Phillipps & Phillipps 2014). They are well known for their important role in regenerating and maintaining tropical rainforest by dispersing seeds (Estrada & Fleming 1986; How 1986; Fleming & Estrada 1993; Hamilton 1999; Hubbell *et al.* 1999; Kinnaird & O'Brien 2007). Apart from regenerating tropical rainforests, hornbills are also useful indicators of forest condition and human disturbance as their habitat preference is for non-fragmented forest and forest with large fruiting trees for their feeding and nesting habit (Gale & Thongaree 2006). They are also known